

1-1-2000

# Choosing a Physical Model: Why Symmetries?

Raul A. Trejo

Vladik Kreinovich

University of Texas at El Paso, vladik@utep.edu

Luc Longpre

University of Texas at El Paso, longpre@utep.edu

Follow this and additional works at: [http://digitalcommons.utep.edu/cs\\_techrep](http://digitalcommons.utep.edu/cs_techrep)

 Part of the [Computer Engineering Commons](#)

Comments:

UTEP-CS-00-04.

Published in *Bulletin of the European Association for Theoretical Computer Science EATCS*, 2000, Vol. 70, pp. 159-161.

---

## Recommended Citation

Trejo, Raul A.; Kreinovich, Vladik; and Longpre, Luc, "Choosing a Physical Model: Why Symmetries?" (2000). *Departmental Technical Reports (CS)*. Paper 460.

[http://digitalcommons.utep.edu/cs\\_techrep/460](http://digitalcommons.utep.edu/cs_techrep/460)

This Article is brought to you for free and open access by the Department of Computer Science at DigitalCommons@UTEP. It has been accepted for inclusion in Departmental Technical Reports (CS) by an authorized administrator of DigitalCommons@UTEP. For more information, please contact [lweber@utep.edu](mailto:lweber@utep.edu).

# Choosing a Physical Model: Why Symmetries?

Raúl A. Trejo, Vladik Kreinovich, and Luc Longpré  
Department of Computer Science  
University of Texas at El Paso  
500 W. University  
El Paso, TX 79968, USA  
{rtrejo,vladik,longpre}@cs.utep.edu

One of the main applications of Kolmogorov complexity ideas to data processing is via the Minimum Description Length principle (see, e.g., [1, 11, 13] and references therein). According to this principle, if several different models (or theories) are consistent with the same observations, then we should choose a model with the shortest description, i.e., crudely speaking, a model with the smallest value of Kolmogorov complexity. This principle is in perfect agreement with the Occam principle (it actually formalizes Occam's principle), and it has been successfully applied to various problems.

In particular, it has been successfully used in physics, where Occam's principle originated and where it has been successfully used. In modern fundamental physics, however, symmetry groups play such an important role that often, physicists choose not the simplest model, but the model which corresponds to the simplest symmetry group (see, e.g., [2–9], [12, 14]).

At first glance, this restriction to *symmetry*-defined models seem to prevent us from considering possible simple *non-group* models, and thus, make this symmetry-group version of Occam principle worse than the unrestricted one. However, the success of this direction in theoretical physics seems to indicate that this restriction does not bring any disadvantage at all. Our analysis shows that this restriction is indeed non-essential.

To formalize the physicists' idea, we fix a universal programming language (“computer”)  $f$ , and define a *symmetry* as a program from this language which transforms strings (e.g., binary strings) into strings and which is a bijection (1-1 and onto). By a *complexity* of a symmetry  $s$ , we mean the length  $\text{len}(s)$  of this program. We say that a symmetry  $s$  *defines a string  $x$  uniquely* if  $s(x) = x$  and  $s(y) \neq y$  for all  $y \neq x$ . Now, for every string  $x$ , we can define its *group-symmetric complexity*  $C_{\text{sym}}(x)$  as the smallest complexity of a symmetry which defines  $x$  uniquely. It turns out that this new complexity is asymptotically equivalent to the usual Kolmogorov complexity  $C(x) = C_f(x)$ :

**Proposition.**  $|C(x) - C_{\text{sym}}(x)| = O(1)$ .

*Comments.*

- Since the Kolmogorov complexity itself is defined modulo the term  $O(1)$ , we conclude that  $C_{\text{sym}}$  is asymptotically equivalent to the usual Kolmogorov complexity. Therefore, choosing the model with the simplest symmetry (i.e., a model  $x$  for which  $C_{\text{sym}}(x)$  is the smallest) is indeed (asymptotically) equivalent to choosing the simplest model.
- This result was first announced in [15].
- An alternative (more physics-oriented) Kolmogorov complexity-based justification of the use of symmetries in physics was given in [10].

**Proof.** To prove that  $|C(x) - C_{\text{sym}}(x)| = O(1)$ , we will first show that  $\forall x(C(x) \leq C_{\text{sym}}(x) + C_1)$  for some constant  $C_1$  and then, that  $\forall x(C_{\text{sym}}(x) \leq C(x) + C_2)$  for some constant  $C_2$ .

Let us prove the first inequality. By definition,  $C_{\text{sym}}(x)$  is the length of a program  $s$  for which  $s(x) = x$  and  $s(y) \neq y$  for all  $y \neq x$ . To compute  $x$ , we can use the following new program  $p$ : Generate, sequentially, all possible binary strings  $\Lambda, 0, 1, 00, 01$ , etc., and check, for each string  $y$ , whether  $s(y) = y$ ; as soon as we get  $s(y) = y$  for some string  $y$ , we stop. This new program  $p$  clearly computes the desired string  $x$ . This program  $p$  has finitely many instructions added to a call to  $s$ ; thus, its length  $\text{len}(p)$  is equal to the length of  $s$  plus the (constant) length  $C_1$  of these extra instructions. Hence,  $\text{len}(p) = \text{len}(s) + C_1 = C_{\text{sym}}(x) + C_1$ . Since the program  $p$  generates  $x$ , the Kolmogorov complexity  $C(x)$  of  $x$  – i.e., the shortest length of the program which generates  $x$  – cannot exceed  $\text{len}(p) = C_{\text{sym}}(x) + C_1$ . Thus, indeed,  $C(x) \leq C_{\text{sym}}(x) + C_1$ . The first inequality is proven.

Let us now prove the second inequality. By definition,  $C(x)$  is the length of a program  $p$  (to be more precise, the shortest program) which generates  $x$ :  $C(x) = \text{len}(p)$ . Based on this program  $p$ , we can construct the desired symmetry  $s(y)$  as follows: For any given input  $y$ , the program  $s$  first calls  $p$  to compute the word  $x$ . Then, the program  $s$  returns the following string  $s(y)$ :

- If  $y = x$ , then we return  $s(x) = x$ .
- If  $y = x^*$ , where by  $x^*$ , we denote a string which differs from  $x$  only in the last bit, then we return the empty string:  $s(x^*) = \Lambda$ .
- If  $y$  is an empty string ( $y = \Lambda$ ), then we return  $s(\Lambda) = x^*$ .
- For all other input strings  $y$ , we return  $s(y) = y^*$ .

One can easily check that this program is indeed a bijection, i.e., a symmetry. For this symmetry, we have  $s(x) = x$  and  $s(y) \neq y$  for all strings  $y \neq x$ ; thus, this symmetry defines the string  $x$  uniquely. The program  $s$  has finitely many instructions added to a call to  $p$ ; hence, its length  $\text{len}(s)$  is equal to the length of  $p$  plus the (constant) length  $C_2$  of these extra instructions. Thus,  $\text{len}(s) = \text{len}(p) + C_2 = C(x) + C_2$ . Since the symmetry  $s$  defines  $x$  uniquely, the group-symmetric complexity  $C_{\text{sym}}(x)$  of  $x$  – i.e., the shortest length of the symmetry which defines  $x$  uniquely – cannot exceed  $\text{len}(s) = C(x) + C_2$ . Thus, indeed,  $C_{\text{sym}}(x) \leq C(x) + C_2$ . The second inequality is also proven, and so is the proposition.

**Acknowledgments.** This work was supported in part by NASA under cooperative agreement NCC5-209, by NSF grants No. DUE-9750858 and CDA-9522207, by United Space Alliance, grant No. NAS 9-20000 (PWO C0C67713A6), by the Future Aerospace Science and Technology Program (FAST) Center for Structural Integrity of Aerospace Systems, effort sponsored by the Air Force Office of Scientific Research, Air Force Materiel Command, USAF, under grant number F49620-95-1-0518, and by the National Security Agency under Grant No. MDA904-98-1-0561.

## References

- [1] C. Calude, *Information and randomness: An algorithmic perspective*, Springer-Verlag, Berlin, 1994.
- [2] A. M. Finkelstein, O. Kosheleva, and V. Kreinovich, “Astrogeometry, error estimation, and other applications of set-valued analysis”, *ACM SIGNUM Newsletter*, 1996, Vol. 31, No. 4, pp. 3–25.
- [3] A. M. Finkelstein, O. Kosheleva, and V. Kreinovich, “Astrogeometry: towards mathematical foundations”, *International Journal of Theoretical Physics*, 1997, Vol. 36, No. 4, pp. 1009–1020.
- [4] A. M. Finkelstein, O. Kosheleva, and V. Kreinovich, “Astrogeometry: geometry explains shapes of celestial bodies”, *Geoinformatics*, 1997, Vol. VI, No. 4, pp. 125–139.
- [5] A. M. Finkelstein and V. Kreinovich, “Derivation of Einstein’s, Brans-Dicke and other equations from group considerations,” *On Relativity Theory. Proceedings of the Sir Arthur Eddington Centenary Symposium, Nagpur India 1984*, Vol. 2, Y. Choque-Bruhat and T. M. Karade (eds), World Scientific, Singapore, 1985, pp. 138–146.
- [6] A. M. Finkelstein, V. Kreinovich, and R. R. Zapatrin, “Fundamental physical equations uniquely determined by their symmetry groups,” *Lecture Notes in Mathematics*, Springer-Verlag, Berlin-Heidelberg-N.Y., Vol. 1214, 1986, pp. 159–170.
- [7] D. R. Finkelstein, *Quantum Relativity: A Synthesis of the Ideas of Einstein and Heisenberg*, Springer-Verlag, Berlin-Heidelberg, 1997.
- [8] *Group theory in physics: proceedings of the international symposium held in honor of Prof. Marcos Moshinsky, Cocoyoc, Morelos, Mexico, 1991*, American Institute of Physics, N.Y., 1992.
- [9] V. Kreinovich, “Derivation of the Schroedinger equations from scale invariance,” *Theoretical and Mathematical Physics*, 1976, Vol. 8, No. 3, pp. 282–285.
- [10] V. Kreinovich and L. Longpré, “Unreasonable effectiveness of symmetry in physics”, *International Journal of Theoretical Physics*, 1996, Vol. 35, No. 7, pp. 1549–1555.
- [11] M. Li and P. Vitányi, *An introduction to Kolmogorov complexity and its applications*, Springer-Verlag, N.Y., 1997.

- [12] P. J. Olver, *Equivalence, invariants, and symmetry*, Cambridge University Press, Cambridge, N.Y., 1995.
- [13] J. J. Rissanen, *Stochastic complexity and statistical inquiry*, World Scientific, 1989.
- [14] *Symmetries in physics: proceedings of the international symposium held in honor of Prof. Marcos Moshinsky, Cocoyoc, Morelos, Mexico, 1991*, Springer-Verlag, Berlin, N.Y., 1992.
- [15] R. A. Trejo, V. Kreinovich, and L. Longpré, “Choosing a Physical Model: Why Symmetries?”, *Complexity Conference Abstracts 1998*, June 1998, Abstract No. 98–37, p. 39.