

From Fuzzy Universal Approximation to Fuzzy Universal Representation: It All Depends on the Continuum Hypothesis

Mahdokht Michelle Afravi and Vladik Kreinovich
Department of Computer Science
University of Texas at El Paso
El Paso, TX 79968, USA
maf ravi@miners.utep.edu, vladik@utep.edu

Abstract

It is known that fuzzy systems have a universal approximation property. A natural question is: can this property be extended to a universal *representation* property? Somewhat surprisingly, the answer to this question depends on whether the following *Continuum Hypothesis*: every infinite subset of the real line has either the same number of elements as the real line itself or as many elements as there are natural numbers.

1 When Can We Go from Fuzzy Universal Approximation to Fuzzy Universal Representation: Formulation of the Problem

Need to translate expert statements into precise terms. In many practical situations, there is a correlation between two quantities x and y , and the only information that we have to describe this correlation are expert statements formulated in terms of imprecise (“fuzzy”) words from natural language, such as “small”.

For example, an expert can say that if x is small, then y is big, and vice versa.

Fuzzy logic provides the desired translation. Fuzzy logic (see, e.g., [4, 5, 6]) is a technique that translates this knowledge into precise mathematical terms.

In this technique, each fuzzy term A is described by a function $A(x)$ assigning, to each possible value x of the corresponding quantity, a degree $A(x)$ to which this value has the appropriate property (e.g., is small).

Once we have rules $A_i(x) \Rightarrow B_i(y)$, the degree $d(x, y)$ to which each pair (x, y) is possible can be described as the degree to which:

- either the first rule is satisfied (i.e., $A_1(x)$ and $B_1(y)$)
- or the second rule is satisfied, etc.

One possible way to interpret “and” is to use product, and “or” is sum. Then, the desired degree takes the form

$$d(x, y) = \sum_{i=1}^n A_i(x) \cdot B_i(y). \quad (1)$$

Universal approximation property. It is known that this expression has a universal approximation property: for every $\varepsilon > 0$, every continuous function on a box can be ε -approximated by such sums.

Is there a universal representation property? A natural question is: when can we get an *exact* representation of every function?

2 Let us Formulate the Problem in Precise Terms

Towards a precise formulation of the problem: first attempt. The simplest way to interpret the above question is to ask whether there exists an integer n such that *any* function of two variables can be represented in the form (1) with this particular n .

It turns out that this is not possible; see, e.g., [2].

Second attempt. Since we cannot have universal representation by using a fixed finite number of terms, a natural next idea is to have a representation in which:

- the number of terms is finite for every function $d(x, y)$ and for every pair (x, y) , but
- this number of terms may be different for different functions $d(x, y)$ and different pairs (x, y) .

Thus, we arrive at the following definition.

Definition 1. *We say that there is a universal representation property if every function $d(x, y)$ of two variables can be represented as the sum*

$$d(x, y) = \sum_{i=1}^{\infty} A_i(x) \cdot B_i(y), \quad (2)$$

so that for every pair (x, y) , only finitely many terms in the sum are different from 0.

Resulting question. So, the resulting question is: is there a universal representation property?

3 Somewhat Surprising Answer: It All Depends on the Continuum Hypothesis

What we should intuitively expect. Intuitively, we expect:

- either a positive answer to the above question – i.e., a proof that the universal representation is possible,
- or a proof that such a universal representation is not possible.

The actual answer is different from our intuition. The actual answer is not what we would intuitively expect.

Proposition 1. [2] *The universal representation property is equivalent to the Continuum Hypothesis.*

Discussion. In a second, we will recall what is the Continuum Hypothesis, but first, let us explain what this result means: that whether we have a universal representation property depends on a somewhat obscure hypothesis from set theory.

What is the Continuum Hypothesis: reminder. In the late 19th century, Georg Cantor invented set theory, the theory which is now the foundations of mathematics. Among other interesting results, he proved that:

- while each infinite subset S of the set N of natural numbers is equivalent to N – in the sense that there is a 1-to-1 correspondence between N and S ,
- the *continuum* – i.e., the set R of real numbers – is *not* equivalent to N (in the above sense).

Cantor conjectured that every infinite subset S of the continuum R is not equivalent either to N or to R . This conjecture became known as *Continuum Hypothesis* (CH). Working mathematicians usually assume this hypothesis, but specialists in foundations of mathematics were interested whether this hypothesis can be proven or disproven based on other – more intuitive – axioms.

This remained an open problem for a long time. The first breakthrough came from the famous logician Kurt Gödel, who proved that the negation of CH cannot be proven in set theory [3]. He proved it by showing that if set theory is consistent, i.e., has a model, then, based on this model, we can build another model in which CH is true.

The question was settled in the 1960s, when Paul Cohen proved that CH is independent of set theory, i.e., we can neither prove nor disprove it based on other axioms of set theory [1]. For this result, he was awarded the Fields Medal – the mathematical equivalent of the Nobel Prize.

Why this result is interesting. At first glance, the Continuum Hypothesis is an obscure statement of set theory, of little interest to working mathematicians

(and probably of even less interest to applications of mathematics). However, surprisingly, this abstract statement is equivalent to something much more practical and interesting: namely, the universal representation property for fuzzy systems.

Of course, one can argue that in practice, when everything is measured and implemented with some accuracy anyway, all we care about is the universal *approximation* property – but still, the universal representation property makes application sense: it shows that we can have an approximation in which:

- for every property $d(x, y)$ and for every pair (x, y) ,
- the number of non-zero terms (i.e., applicable expert rules) remains constant no matter how much accuracy we seek.

Acknowledgments

This work was supported in part by the National Science Foundation grants HRD-0734825 and HRD-1242122 (Cyber-ShARE Center of Excellence) and DUE-0926721, and by an award “UTEP and Prudential Actuarial Science Academy and Pipeline Initiative” from Prudential Foundation.

References

- [1] P. K. Cohen, *Set theory and the continuum hypothesis*, W. A. Benjamin, New York, 1966.
- [2] R. O. Davies, “Representation of functions of two variables as sums of rectangular functions I”, *Fundamenta Mathematicae*, 1974, Vol. 85, No. 2, pp. 177–183.
- [3] K. Gödel, *The Consistency of the Continuum-Hypothesis*, Princeton University Press, Princeton, New Jersey, 1940.
- [4] G. Klir and B. Yuan, *Fuzzy Sets and Fuzzy Logic*, Prentice Hall, Upper Saddle River, New Jersey, 1995.
- [5] H. T. Nguyen and E. A. Walker, *A First Course in Fuzzy Logic*, Chapman and Hall/CRC, Boca Raton, Florida, 2006.
- [6] L. A. Zadeh, “Fuzzy sets”, *Information and Control*, 1965, Vol. 8, pp. 338–353.